

MORTALITY FROM SMOKING IN DEVELOPED COUNTRIES 1950–2020

(see also www.deathsfromsmoking.net)

**Richard Peto, Alan D Lopez, Hongchao Pan
Jillian Boreham and Michael Thun**

MALTA

Raw data (pages 310–11)

Analyses (pages 312–13, 314–15, 316–17)

Appendix (pages 318–19)

(revised September 2015)

MALTA: 2010

	No. of deaths (Source: WHO)			Standardised rates (defined overleaf)			Annual death rates / 100,000 (zero denoted by -)						
	All ages	0-34	35-69	All ages	0-34	35-69	0-4	5-9	10-14	15-19	20-24	25-29	30-34
ALL CAUSES	M 1489	54	473	607.1	58.8	530.8	149.6	-	16.4	37.6	60.9	39.5	107.5
	F 1521	22	281	424.5	29.9	292.3	116.7	-	22.1	7.1	29.5	27.9	6.2
Tuberculosis	M	-	-	-	-	-	-	-	-	-	-	-	-
	F	1	-	0.4	-	-	-	-	-	-	-	-	-
Other infective and parasitic	M	1	-	1	0.5	-	1.0	-	-	-	-	-	-
	F	3	-	0.9	-	-	-	-	-	-	-	-	-
ALL CANCER	M 455	5	215	179.0	5.2	241.1	-	-	16.4	7.5	-	-	12.6
	F 397	3	170	130.2	4.0	175.4	10.6	-	7.4	-	9.8	-	-
Mouth and pharynx cancer	M	12	-	8	5.3	-	9.1	-	-	-	-	-	-
	F	7	-	3	2.2	-	3.1	-	-	-	-	-	-
Oesophagus cancer	M	5	-	-	1.8	-	-	-	-	-	-	-	-
	F	4	-	-	1.1	-	-	-	-	-	-	-	-
Stomach cancer	M	22	-	9	8.5	-	11.2	-	-	-	-	-	-
	F	9	-	1	2.8	-	1.0	-	-	-	-	-	-
Colorectal cancer	M	60	-	28	23.5	-	31.8	-	-	-	-	-	-
	F	51	-	22	15.6	-	23.6	-	-	-	-	-	-
Liver cancer	M	8	-	6	3.1	-	6.3	-	-	-	-	-	-
	F	4	-	1	1.3	-	0.9	-	-	-	-	-	-
Pancreas cancer	M	41	-	23	16.1	-	25.3	-	-	-	-	-	-
	F	32	-	14	11.2	-	14.4	-	-	-	-	-	-
Larynx cancer	M	4	-	3	1.5	-	3.0	-	-	-	-	-	-
	F	1	-	-	0.3	-	-	-	-	-	-	-	-
Lung cancer	M	122	-	62	47.1	-	69.7	-	-	-	-	-	-
	F	37	-	19	12.6	-	20.0	-	-	-	-	-	-
Malignant melanoma	M	3	-	1	1.3	-	1.4	-	-	-	-	-	-
	F	3	-	1	1.2	-	1.0	-	-	-	-	-	-
Female breast cancer	F	79	-	37	26.6	-	37.9	-	-	-	-	-	-
Cervix cancer	F	2	-	-	0.6	-	-	-	-	-	-	-	-
Other uterine cancer	F	17	-	8	6.0	-	8.0	-	-	-	-	-	-
Ovarian cancer	F	37	-	21	13.3	-	21.5	-	-	-	-	-	-
Prostate cancer	M	30	-	4	11.5	-	4.3	-	-	-	-	-	-
Bladder cancer	M	29	-	9	11.4	-	9.6	-	-	-	-	-	-
	F	4	-	-	1.1	-	-	-	-	-	-	-	-
Other and ill-defined cancer sites	M	81	1	42	32.2	1.2	46.5	-	-	8.2	-	-	-
	F	76	2	30	24.0	2.5	30.3	-	-	7.4	-	9.8	-
Hodgkin's disease	M	1	-	1	0.3	-	0.9	-	-	-	-	-	-
	F	2	-	-	0.6	-	-	-	-	-	-	-	-
Myeloma and non-Hodgkin lymphomas	M	17	1	10	6.9	0.9	11.3	-	-	-	-	-	6.3
	F	18	-	8	5.2	-	8.6	-	-	-	-	-	-
Leukaemia	M	20	3	9	8.5	3.1	10.6	-	-	8.2	7.5	-	6.3
	F	14	1	5	4.6	1.5	4.9	10.6	-	-	-	-	-
Diabetes	M	47	-	18	18.8	-	20.3	-	-	-	-	-	-
	F	56	-	7	15.0	-	7.3	-	-	-	-	-	-
ALL VASCULAR DISEASE	M 510	2	135	206.8	1.8	151.4	-	-	-	-	-	-	12.6
Rheumatic heart disease and fever	M	642	3	54	163.9	3.3	56.4	-	-	7.4	-	-	9.3
	F	5	-	1	0.7	-	1.4	-	-	-	-	-	6.2
Hypertensive disease	M	7	-	2	3.0	-	2.4	-	-	-	-	-	-
	F	26	-	3	5.8	-	3.3	-	-	-	-	-	-
Ischaemic heart disease	M	319	1	91	129.3	0.9	101.3	-	-	-	-	-	6.3
	F	327	-	23	85.0	-	24.5	-	-	-	-	-	-
Pulmonary embolism and other venous	M	5	-	1	2.1	-	1.0	-	-	-	-	-	-
	F	10	1	1	3.0	1.3	0.9	-	-	-	-	9.3	-
Cerebrovascular disease	M	103	1	20	41.4	0.9	22.6	-	-	-	-	-	6.3
	F	161	1	12	40.0	1.1	12.2	-	-	7.4	-	-	-
Other vascular disease	M	74	-	20	30.4	-	22.8	-	-	-	-	-	-
	F	113	1	14	28.6	0.9	14.3	-	-	-	-	-	6.2
Chronic obstructive pulmonary disease	M	72	1	8	29.0	1.1	9.1	-	-	7.5	-	-	-
	F	18	-	1	4.6	-	1.0	-	-	-	-	-	-
Other respiratory disease	M	92	-	10	37.2	-	11.8	-	-	-	-	-	-
	F	111	-	10	26.8	-	10.4	-	-	-	-	-	-
Peptic ulcer	M	2	-	1	0.9	-	1.0	-	-	-	-	-	-
	F	3	-	-	0.8	-	-	-	-	-	-	-	-
Liver cirrhosis	M	16	-	11	6.2	-	13.0	-	-	-	-	-	-
	F	4	-	2	1.4	-	2.2	-	-	-	-	-	-
Renal disease	M	22	-	3	8.9	-	2.9	-	-	-	-	-	-
	F	28	-	4	7.1	-	4.5	-	-	-	-	-	-
Pregnancy, birth	F	-	-	-	-	-	-	-	-	-	-	-	-
Congenital and perinatal causes	M	14	12	-	10.0	16.6	-	109.7	-	-	6.8	-	-
	F	13	10	1	9.5	15.2	0.9	106.1	-	-	-	-	-
Ill-defined causes	M	17	2	4	7.2	1.9	4.1	-	-	-	6.8	6.6	-
	F	16	-	1	3.5	-	1.1	-	-	-	-	-	-
Other medical causes	M	152	9	26	62.2	10.3	30.6	39.9	-	-	13.5	-	19.0
	F	199	-	23	49.8	-	24.5	-	-	-	-	-	-
ALL NON-MEDICAL CAUSES	M 89	23	41	40.5	21.8	44.5	-	-	22.6	33.9	32.9	63.2	-
Road traffic accidents	M	30	6	8	10.5	7.5	8.4	-	-	7.4	7.1	19.7	18.6
	F	11	6	3	5.0	5.7	3.4	-	-	-	7.5	6.8	25.3
Fire	M	1	-	-	0.4	-	-	-	-	-	-	-	-
	F	-	-	-	-	-	-	-	-	-	-	-	-
Suicide	M	29	6	20	14.1	5.9	21.4	-	-	15.1	6.8	6.6	12.6
	F	2	-	2	0.7	-	2.1	-	-	-	-	-	-
POPULATION: (1000s; UN, 2015 revision)	M=male	204.7	91.9	94.2	10.0	10.6	12.2	13.3	14.8	15.2	15.8		
	F=female	207.4	85.6	96.7	9.4	11.3	13.6	14.1	10.2	10.8	16.3		

2010: MALTA

Annual death rates / 100,000 (zero denoted by -)

9th ICD/10th ICD categories

35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80+/NK					
101.2	178.0	178.0	359.7	597.4	929.4	1372.0	2078.6	4902.9	11102.2	M	ALL CAUSES	001-999/A00-Y89		
21.1	60.8	136.1	211.5	288.2	606.8	721.6	1459.2	4315.2	8601.2	F	Tuberculosis	010-018, 137/ A15-A19, B90		
-	-	-	-	-	-	-	-	-	-	M	Other infective and parasitic	Rest of 001-139, 279, 5&6/ rest of A00-B99 excl A33,A34		
7.2	-	-	-	-	-	-	-	-	37.5	10.2	F			
7.2	23.2	54.8	148.1	280.2	517.8	656.2	897.1	1310.4	1900.4	M	ALL CANCER	140-208/C00-C97		
21.1	30.4	93.1	155.1	184.7	382.5	360.8	533.4	1275.8	1050.9	F				
-	15.5	6.8	14.1	7.4	-	19.9	21.9	22.6	20.0	M	Mouth and pharynx cancer	140-149/C00-C14		
-	-	-	-	7.4	6.6	8.0	-	37.5	20.4	F	Oesophagus cancer	150/C15		
-	-	-	-	-	-	-	10.1	18.8	20.4	M				
-	-	-	-	-	-	-	-	-	-	F				
-	-	-	-	22.1	6.6	49.7	65.6	45.2	100.0	M	Stomach cancer	151/C16		
-	-	7.2	-	-	-	-	10.1	56.3	40.8	F				
-	-	6.8	7.0	29.5	79.7	99.4	65.6	180.8	360.1	M	Colorectal cancer	153, 154/C18-C21		
-	-	-	14.1	29.6	33.0	88.2	40.2	150.1	173.4	F				
-	-	-	7.0	7.4	19.9	9.9	10.9	-	20.0	M	Liver cancer	155.0/C22 excl C221		
-	-	-	-	-	6.6	-	-	37.5	10.2	F				
-	7.7	-	14.1	29.5	66.4	59.6	54.7	113.0	160.0	M	Pancreas cancer	157/C25		
-	7.6	7.2	21.1	14.8	26.4	24.1	30.2	168.9	61.2	F				
-	-	-	-	7.4	13.3	-	-	-	20.0	M	Larynx cancer	161/C32		
-	-	-	-	-	-	-	10.1	-	-	F				
-	-	13.7	49.4	66.4	159.3	198.8	306.3	361.5	320.1	M	Lung cancer	162/C33, C34		
-	15.2	-	21.1	14.8	33.0	56.1	60.4	112.6	61.2	F				
-	-	-	-	-	-	9.9	-	45.2	-	M	Malignant melanoma	172/C43		
-	-	-	7.0	-	-	-	10.1	18.8	-	F				
7.0	-	14.3	42.3	51.7	85.7	64.1	110.7	281.4	163.2	F	Female breast cancer	174/C50		
-	-	-	-	-	-	-	20.1	-	-	F	Cervix cancer	180/C53		
-	-	-	14.1	7.4	26.4	8.0	40.2	75.0	10.2	F	Other uterine cancer	179, 182/C54, C55		
7.0	-	28.6	21.1	22.2	39.6	32.1	80.5	75.0	40.8	F	Ovarian cancer	183/C56, C570-C574		
-	-	-	7.0	-	13.3	9.9	98.5	67.8	280.1	M	Prostate Cancer	185/C61		
-	-	-	7.0	7.4	33.2	19.9	43.8	135.6	200.0	M	Bladder cancer	188/C67		
-	-	-	-	-	-	-	10.1	18.8	20.4	F				
7.2	-	27.4	28.2	73.8	79.7	109.4	175.0	158.2	300.1	M	Other and ill-defined cancer sites	Rest of 140-199/ rest of C00-C80, C97		
-	7.6	21.5	7.0	37.0	98.9	40.1	80.5	187.6	265.3	F	Hodgkin's disease	201/C81		
-	-	-	-	-	6.6	-	-	-	18.8	10.2	M			
-	-	-	7.0	22.1	19.9	29.8	21.9	67.8	20.0	M	Myeloma and non- Hodgkin lymphomas	200, 202-203/ C82-C90, C96		
7.0	-	-	-	-	13.2	40.1	20.1	18.8	71.4	F				
-	-	-	7.0	7.4	19.9	39.8	-	113.0	60.0	M	Leukaemia	204-208/C91-C95		
-	-	14.3	7.0	-	13.2	-	-	-	81.6	F				
-	-	6.8	21.2	14.8	39.8	59.6	65.6	158.2	320.1	M	Diabetes	250/E10-E14		
-	-	-	-	7.4	19.8	24.1	60.4	225.1	316.3	F				
7.2	54.2	68.5	127.0	206.5	219.1	377.8	656.4	2010.8	4480.9	M	ALL VASCULAR DISEASE	390-459/I00-I99		
-	15.2	14.3	35.2	59.1	118.7	152.3	533.4	1932.5	4377.1	F	Rheumatic heart disease and fever	390-398/I00-I09		
-	-	-	-	-	-	9.9	10.9	-	-	M	Hypertensive disease	401-405/I10-I15		
-	-	-	6.8	-	-	8.0	-	37.5	20.4	F				
-	-	-	7.2	-	-	9.9	-	45.2	60.0	M				
-	-	-	-	-	16.0	-	-	-	234.7	F				
7.2	15.5	41.1	91.7	162.2	152.7	238.6	426.6	1310.4	2600.5	M	Ischaemic heart disease	410-414/I20-I25		
-	7.6	7.2	21.1	22.2	33.0	80.2	301.9	1238.3	2122.2	F	Pulmonary embolism	415.1, 451-453/ I26, I80-I82		
-	-	6.8	-	-	-	-	10.9	45.2	21.0	M	and other venous	430-438/I60-I69		
-	-	-	-	-	6.6	-	10.1	37.5	51.0	F	Cerebrovascular disease			
-	7.7	-	14.1	36.9	39.8	59.6	142.2	429.3	1000.2	M				
-	-	-	7.0	14.8	39.6	24.1	120.8	412.8	1163.2	F				
-	-	30.9	13.7	21.2	7.4	26.6	65.6	180.8	800.2	M	Other vascular disease	Rest of 390-459/ rest of I00-I99		
-	7.6	-	7.0	22.2	39.6	24.1	100.6	206.4	785.6	F				
-	-	-	-	7.4	26.6	29.8	98.5	429.3	700.1	M	Chronic obstructive pulmonary disease	490-496/ J40-J47, J67		
-	-	-	7.0	-	-	-	30.2	37.5	122.4	F	Other respiratory disease	Rest of 460-519/ rest of J00-J98		
-	-	15.5	-	7.0	7.4	13.3	39.8	65.6	248.5	1300.3	M			
-	-	-	7.0	7.4	26.4	32.1	100.6	187.6	826.5	F				
-	-	-	-	7.0	-	-	-	-	-	M	Peptic ulcer	531-533/K25-K27		
-	-	-	-	-	-	-	-	-	20.0	F				
-	-	6.8	7.0	7.4	19.9	49.7	43.8	22.6	-	20.4	F	Liver cirrhosis	571/K70, K74	
-	-	-	-	7.4	-	8.0	10.1	18.8	-	M				
-	-	-	-	-	19.9	-	21.9	203.3	160.0	M	Renal disease	580-590/N00-N19		
-	-	-	-	7.4	-	24.1	10.1	75.0	193.9	F				
-	-	-	-	-	-	-	-	-	-	F	Pregnancy, birth Congenital and perinatal causes	630-676/A34,000-O99 740-779/A33, P00-P96, Q00-Q99		
-	-	-	-	-	6.6	-	10.9	-	-	M	III-defined causes	780-799/R00-R99		
-	-	-	7.0	14.8	6.6	-	-	45.2	180.0	M				
-	-	-	-	-	-	8.0	-	18.8	142.8	F				
-	-	15.5	6.8	-	29.5	53.1	109.4	186.0	361.5	1680.3	M	Other medical causes	Rest of 001-799/ rest of A00-R99	
-	7.6	21.5	-	14.8	39.6	88.2	150.9	431.5	1408.0	F				
79.5	69.7	34.2	35.3	29.5	13.3	49.7	32.8	113.0	340.1	M	ALL NON- MEDICAL CAUSES	E800-E999/V01-Y89		
-	7.6	7.2	7.0	-	13.2	24.1	20.1	18.8	132.6	F	Road traffic accidents	E810-E819, E826-E829/ part of V01-V99, Y850		
-	-	6.8	-	-	-	9.9	-	22.6	20.0	M				
-	-	-	-	-	-	-	10.1	-	-	F				
-	-	36.2	61.9	6.8	21.2	7.4	6.6	9.9	10.9	22.6	20.0	M	Fire	E890-E899/X00-X09
-	-	-	-	-	-	-	-	-	-	-	F	Suicide	E950-E959/X60-X84	
13.8	12.9	14.6	14.2	13.6	15.1	10.1	9.1	4.4	5.0	M	POPULATION (1000s; UN, 2015 revision)			
14.2	13.2	14.0	14.2	13.5	15.2	12.5	9.9	5.3	9.8	F				

MALTA: 2010**Relative importance of deaths in MIDDLE age (35–69) in the year 2010**

Age range (years)	Deaths attributed to SMOKING /total deaths		Mean years lost PER DEATH FROM SMOKING
	Male	Female	
0–34	– / 56	– / 31	–
35–69	131 / 501	11 / 293	22 years
70+	256 / 1093	63 / 1261	7 years
All ages	387 / 1650	74 / 1585	12 years



Deaths, by cause, attributed to SMOKING / total deaths in the year 2010

Cause	Male (by age)				Female (by age)			
	0–34	35–69	70+	All	0–34	35–69	70+	All
Lung Cancer	–/0	47/54	67/74	114/128	–/0	5/15	8/15	13/30
All Cancer	–/6	65/198 (33%)	94/266 (35%)	159/470	–/4	5/162 (3%)	13/225 (6%)	18/391
Vascular	–/2	47/170	91/483	138/655	–/3	5/67	32/621	37/691
Respiratory	–/0	10/21	52/133	62/154	–/0	0/10	10/114	10/124
All Other	–/43	9/110	19/210	28/363	–/22	1/51	8/300	9/373
All Causes	–/56	131/501 (26%)	256/1093 (23%)	387/1650	–/31	11/293 (4%)	63/1261 (5%)	74/1585

**Cancer deaths, and all deaths,
attributed to SMOKING / total deaths in the year 2010**



Cause	Male	Female	Male + Female
All Cancer	159 / 470 (34%)	18 / 391 (5%)	177 / 861 (21%)
All Causes	387 / 1650 (23%)	74 / 1585 (5%)	461 / 3235 (14%)

1965-2015: MALTA

Population risk of dying at ages 0–34**Population risk of a 35-year-old dying at ages 35–69 from smoking (shaded) or from any cause (shaded and white)****MALE**

*eg, at year 2010 male death rates, out of 100 men aged 35, 18 would die before age 70 (with 5 of these deaths attributed to smoking)

Note: Most of those killed by smoking would otherwise have survived beyond age 70, but a minority (shaded area to right of dotted line) would have died by 70 anyway

FEMALE

Real risk too low to estimate reliably

MALTA: 1960-2015***Relative importance of deaths in MIDDLE age (35–69), 2015 projections***

Age range (years)	Deaths attributed to SMOKING /total deaths		Mean years lost PER DEATH FROM SMOKING
	Male	Female	
0–34	– / 57	– / 44	–
35–69	125 / 486	31 / 268	24 years
70+	323 / 1175	120 / 1268	7 years
All ages	448 / 1718	151 / 1580	11 years

Numbers of deaths attributed to smoking / total deaths

Year	Male (by age)				Female (by age)			
	0–34	35–69	70+	All	0–34	35–69	70+	All
1955
1965	–/165	237/631 (38%)	67/747 (9%)	304/1543	–/117	0/486	0/835	0/1438
1975	–/113	235/616 (38%)	145/806 (18%)	380/1535	–/81	1/426	0/923	1/1430
1985	–/94	226/540 (42%)	188/891 (21%)	414/1525	–/58	0/361	0/1029	0/1448
1995	–/75	174/451 (39%)	234/901 (26%)	408/1427	–/37	0/305	0/981	0/1323
2005	–/56	124/466 (27%)	233/1065 (22%)	357/1587	–/26	4/301	30/1208	34/1535
2015 proj.	–/57	125/486 (26%)	323/1175 (27%)	448/1718	–/44	31/268	120/1268 (9%)	151/1580





60-year total (thousands), decade of 1960s to decade of 2010s (1960–2020*): 25 / 181

*Estimated as 10 times the sum of the annual numbers for 1965, 1975, 1985, 1995, 2005 and 2015

1960–2020, by age & sex:


–/5.6	11/32 (35%)	12/56 (21%)	23/93	–/3.6	0.4/21 (2%)	1.5/62 (2%)	1.9/88
-------	-----------------------	-----------------------	-------	-------	-----------------------	-----------------------	--------

1965-2015: MALTA



MALTA: 1965-2015

ALL-CAUSE mortality rates attributed and not attributed to smoking




Year	Attributed to smoking?	MALE all-cause mortality (annual rate/1000, by age)						FEMALE all-cause mortality (annual rate/1000, by age)					
		0–34		35–69		70–79		0–34		35–69		70–79	
		Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No
1950	
1955	
1960	
1965	-	1.61	6.29	11.0	9.75	84.2	-	1.22	0.00	12.9	0.00	75.9	
1970	-	1.51	6.88	10.2	14.0	73.5	-	1.09	0.00	10.9	0.00	77.0	
1975	-	1.26	5.93	9.92	15.9	66.8	-	0.91	0.02	9.28	0.00	73.8	
1980	-	1.07	5.96	8.79	16.9	71.7	-	0.71	0.00	9.07	0.00	69.3	
1985	-	0.94	5.12	7.29	18.3	59.5	-	0.60	0.00	7.25	0.00	53.0	
1990	-	0.79	3.85	6.19	16.1	48.9	-	0.44	0.00	5.95	0.00	43.1	
1995	-	0.78	3.24	5.02	16.5	43.2	-	0.40	0.00	5.09	0.00	38.0	
2000	-	0.64	2.72	5.11	13.1	43.7	-	0.33	0.12	4.28	1.51	34.1	
2005	-	0.61	1.91	4.75	12.4	35.1	-	0.33	0.05	3.49	0.79	27.9	
2010	-	0.62	1.50	4.22	10.2	28.9	-	0.41	0.11	2.97	1.29	25.8	
2015	-	0.65	1.29	3.70	9.84	23.5	-	0.61	0.32	2.42	2.13	24.6	

0–34 and 35–69 = means of 7 age-specific rates; 70–79 = mean of 2 rates (70–74 & 75–79)


1965-2015: MALTA

ALL CANCER mortality rates attributed and not attributed to smoking

[†] Mean of 7 age-specific rates, ages 35–69 (or ages 0–34)

Year	MALE cancer mortality (annual rate/1000, by age)						FEMALE cancer mortality (annual rate/1000, by age)								
	Attributed to smoking?	0–34		35–69		70–79		Yes	No	0–34		35–69		70–79	
		Yes	No	Yes	No	Yes	No			Yes	No	Yes	No	Yes	No
1950		
1955		
1960		
1965	-	0.06	1.46	2.24	1.82	7.58	-	0.11	0.00	2.21	0.00	7.62	-	0.09	
1970	-	0.11	1.50	1.83	2.75	6.87	-	0.09	0.00	2.26	0.00	7.72	-	0.10	
1975	-	0.09	1.36	1.82	3.46	6.92	-	0.07	0.00	1.95	0.00	7.24	-	0.09	
1980	-	0.07	1.60	1.85	4.15	8.90	-	0.09	0.00	2.32	0.00	8.64	-	0.08	
1985	-	0.10	1.58	1.67	5.32	8.81	-	0.07	0.00	2.23	0.00	8.17	-	0.07	
1990	-	0.05	1.41	1.78	5.15	9.27	-	0.07	0.00	2.09	0.00	7.82	-	0.06	
1995	-	0.09	1.41	1.73	5.79	10.4	-	0.03	0.00	2.27	0.00	8.55	-	0.05	
2000	-	0.06	1.15	1.76	4.74	10.1	-	0.05	0.03	1.95	0.30	7.56	-	0.04	
2005	-	0.06	0.89	1.53	4.69	8.59	-	0.05	0.02	1.79	0.22	7.01	-	0.04	
2010	-	0.06	0.75	1.52	4.66	8.04	-	0.05	0.05	1.65	0.39	7.12	-	0.04	
2015	-	0.07	0.74	1.46	5.17	7.26	-	0.16	0.16	1.47	0.73	7.59	-	0.05	


0–34 and 35–69 = means of 7 age-specific rates; 70–79 = mean of 2 rates (70–74 & 75–79)

MALTA: 1965-2012**Mortality* trends at selected ages: 35-39, 55-59 & 75-79**

* Annual mortality per 1000: averages of years available in 1965-69,...,2005-09; then 3-year average 2010-12

1965-2012: MALTA

Mortality* trends at selected ages: 35-39, 55-59 & 75-79



* Annual mortality per 1000: averages of years available in 1965-69,...,2005-09; then 3-year average 2010-12